

VETRINARIAN

Lipoic Acid Mineral Complex

Chronic / Neurological / Cancer / Palliative / General Care

Veterinarian Frequency/ Options for IM/Subcutaneous- IV Protocols

- a. 1 or 4 days per week for 4-12 consecutive weeks
- b. Intermittently to coincide with or around other IV protocols or treatments

Case studies have shown to titrate on oral Poly-MVA before beginning the IV protocols.

The patient will then be doing both the IV and oral Poly simultaneously. The oral dose is between 1-4 tsp 5-20ml per day (best for divided doses) you can mix Poly-MVA with a favorite beverage (water, juice, or broth etc.).

Maintain the oral dose for at least 2-3 months or as needed.

The minimum effective dose for continuing patients, depending on body weight is at least 1/2 - 2 tsp/ day.

Subcutaneous: 3-10cc using subcue, you can dilute with saline if necessary.

IV preparation and administration:

1. **Generally start with 50-100 cc normal saline.** Initially add 5-15 cc of injectable LAMC.
Increase each infusion by 5cc of LAMC until max of 40 cc or 0.5 cc per kg (based on an 80 kg patient) or 1-2ml per 5lbs of body weight.
2. Run the infusions over 30-45 minutes.
3. 10-20cc 3 times for first week with oral dosage of 10-20 cc on the off days.
4. Dosage can increase to 30/40 cc of LAMC, **increase to 250/500 cc of saline if desired.**
Run this over one hour or more.
5. 30cc 3 times for second week with oral dosage of 10cc
6. 40cc 3 times for third week with oral on days off and weekends.

On days with no IV Poly, use oral LAMC between 5-40ml per day.